

Struggling with unresolved depression symptoms?
Ask about adding VRAYLAR to your antidepressant.

FOR A LIFT IN RELIEF



Vraylar[®]
(cariprazine) capsules
1.5mg•3mg

VRAYLAR is a prescription medicine used along with antidepressants to treat major depressive disorder (MDD) in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VRAYLAR?

Elderly people with dementia-related psychosis (having lost touch with reality due to confusion and memory loss) taking medicines like VRAYLAR are at an increased risk of death. VRAYLAR is not approved for treating patients with dementia-related psychosis.

VRAYLAR and antidepressants may increase suicidal thoughts or actions in some children and young adults especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. This is very important when VRAYLAR or the antidepressant is started or when the dose is changed. Report any change in these symptoms immediately to the doctor.

Please see Important Safety Information throughout and full [Prescribing Information](#), including [Boxed Warnings](#), and [Medication Guide](#).

Could VRAYLAR be right for me?



If you have unresolved depression symptoms even on your antidepressant, **ask your healthcare provider today if VRAYLAR could be right for you.**

IMPORTANT SAFETY INFORMATION (continued)

VRAYLAR may cause serious side effects, including:

- **Stroke (cerebrovascular problems)** in elderly people with dementia-related psychosis that can lead to death
- **Neuroleptic malignant syndrome (NMS):** Call your healthcare provider or go to the nearest hospital emergency room right away if you have high fever, stiff muscles, confusion, increased sweating, or changes in breathing, heart rate, and blood pressure. These can be symptoms of a rare but potentially fatal side effect called NMS. VRAYLAR should be stopped if you have NMS.
- **Uncontrolled body movements (tardive dyskinesia or TD):** VRAYLAR may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking VRAYLAR. Tardive dyskinesia may also start after you stop taking VRAYLAR.
- **Late-occurring side effects:** VRAYLAR stays in your body for a long time. Some side effects may not happen right away and can start a few weeks after starting VRAYLAR, or if your dose increases. Your healthcare provider should monitor you for side effects for several weeks after starting or increasing dose of VRAYLAR.



What is VRAYLAR?

VRAYLAR is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

VRAYLAR is a once-daily pill that can be taken with or without food.



How could VRAYLAR help me?

When added to an antidepressant, VRAYLAR is clinically proven to help relieve **overall depression symptoms** better than an antidepressant alone.

Individual results may vary.



How does VRAYLAR work?

VRAYLAR works with your antidepressant. VRAYLAR also **works differently than an antidepressant**. While the exact way VRAYLAR works is unknown, scientists and doctors believe that VRAYLAR helps manage chemicals in the brain, such as serotonin and dopamine, that may control your mood.



Do I need to change my current treatment?

No. If you've made progress on your antidepressant but are still experiencing unresolved depression symptoms, your doctor may add VRAYLAR to help give your antidepressant a lift.

Please see Important Safety Information throughout and full [Prescribing Information](#), including [Boxed Warnings](#), and [Medication Guide](#).



What antidepressants could I take with VRAYLAR?

In VRAYLAR clinical studies, patients were taking commonly used antidepressants,* including:

- Fluoxetine (Prozac®)
- Venlafaxine (Effexor®)
- Paroxetine (Paxil®)
- Bupropion hydrochloride (Wellbutrin®)
- Sertraline (Zoloft®)
- Escitalopram (Lexapro®)
- Duloxetine (Cymbalta®)
- Citalopram (Celexa®)


These are not all the antidepressants studied with VRAYLAR. Ask your healthcare provider if adding VRAYLAR is right for you.

*All trademarks are property of their respective owners.

Will I gain weight while on VRAYLAR?

Most people taking VRAYLAR and an antidepressant (ADT) saw no substantial impact on weight.

Average weight change in 6-week major depressive disorder studies:



In clinical studies, 98% of patients in 6-week trials and 81% of patients in a 6-month trial had no substantial weight gain (defined by $\geq 7\%$ change). In the 6-month study, the average weight gain in VRAYLAR-treated patients was 3.7 lb. Weight gain may occur. You and your doctor should monitor your weight regularly.

IMPORTANT SAFETY INFORMATION (continued)

VRAYLAR may cause serious side effects, including:

- Problems with your metabolism, such as:
 - **High blood sugar and diabetes:** Increases in blood sugar can happen in some people who take VRAYLAR. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before or soon after starting VRAYLAR and regularly during treatment. Tell your healthcare provider if you have symptoms such as feeling very thirsty, very hungry, or sick to your stomach, urinating more than usual, feeling weak, tired, confused, or your breath smells fruity.
 - **Increased fat levels (cholesterol and triglycerides) in your blood:** Your healthcare provider should check fat levels in your blood before or soon after starting VRAYLAR and during treatment.



A lift in relief is possible with VRAYLAR
Talk to your healthcare provider about adding
VRAYLAR to your current antidepressant.

Can I anticipate any sexual side effects when adding VRAYLAR?

In clinical studies, less than 1% of patients taking VRAYLAR with their antidepressant reported sexual side effects.*

	Placebo + ADT	VRAYLAR 1.5 mg/day + ADT	VRAYLAR 3 mg/day + ADT
Abnormal Orgasm	0%	0%	0.2%
Decreased Sex Drive	0%	0.2%	0%
Erectile Dysfunction	0%	0.2%	0%
Ejaculation Failure	0%	0.2%	0%
Ejaculation Disorder	0%	0%	0%

*These adverse events were self-reported by patients in 6-week studies.

IMPORTANT SAFETY INFORMATION (continued)

VRAYLAR may cause serious side effects, including:

- **Problems with your metabolism, such as (continued):**
 - **Weight gain:** Weight gain has been reported with VRAYLAR. You and your healthcare provider should check your weight before and regularly during treatment.
- **Low white blood cell count:** Low white blood cell counts have been reported with antipsychotic drugs, including VRAYLAR. This may increase your risk of infection. Very low white blood cell counts, which can be fatal, have been reported with other antipsychotics. Your healthcare provider may do blood tests during the first few months of treatment with VRAYLAR.

Please see Important Safety Information throughout and full Prescribing Information, including Boxed Warnings, and Medication Guide.

**You've made progress with your antidepressant.
Adding VRAYLAR may help get you where you want to be.**



IMPORTANT SAFETY INFORMATION (continued)


VRAYLAR may cause serious side effects, including:

- **Decreased blood pressure (orthostatic hypotension):** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls:** VRAYLAR may make you sleepy or dizzy, may cause a decrease in blood pressure when changing position (orthostatic hypotension), and can slow thinking and motor skills, which may lead to falls that can cause fractures or other injuries.
- **Seizures (convulsions)**
- **Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities:** Do NOT drive, operate machinery, or do other dangerous activities until you know how VRAYLAR affects you. VRAYLAR may make you drowsy.
- **Increased body temperature:** Do not become too hot or dehydrated during VRAYLAR treatment. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing** that can cause food or liquid to get into your lungs

Learn how you could save.

Join our savings program, VRAYPAYSM, and you may be able to decrease your out-of-pocket costs.*

To enroll, you can:



Scan this **QR code**

TEXT **SAVE TO 20686**[†]

Go to [VRAYLAR.com/cost-and-savings](https://vraylar.com/cost-and-savings)

Not Actual Card Image

* Eligibility: Available to patients with commercial insurance coverage for VRAYLAR[®] (cariprazine) who meet eligibility criteria. This copay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare, Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. For full Terms and Conditions, visit vraypay.com or call 1-800-761-0436 for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit <https://privacy.abbvie/>

[†] Text SAVE to 20686 to enroll or activate your savings card and receive refill reminders. Message and data rates may apply. Message frequency varies. You are not required to consent as a condition of receiving goods or services. You can reply HELP for help. You can reply STOP to opt out at any time. By texting SAVE to 20686, you agree to program Terms and Conditions, AbbVie Privacy Policy, and VRAYLAR SMS Terms.

IMPORTANT SAFETY INFORMATION (continued)

Who should not take VRAYLAR?

Do not take VRAYLAR if you are allergic to any of its ingredients. Get emergency medical help if you are having an allergic reaction (eg, rash, itching, hives, swelling of the tongue, lip, face or throat).

What should I tell my healthcare provider before taking VRAYLAR?

Tell your healthcare provider about any medical conditions and if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar in you or your family
- have or have had high levels of total cholesterol, LDL-cholesterol, or triglycerides; or low levels of HDL-cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had low white blood cell count

Please see Important Safety Information throughout and full Prescribing Information, including Boxed Warnings, and Medication Guide.

Vraylar[®]

(cariprazine) capsules
1.5mg-3mg

Through our work with advocacy groups and patients, AbbVie, the maker of VRAYLAR, is helping to foster discussion on mental health and reduce the stigma around it.

Visit [VraylarMDD.com](https://www.vraylarmdd.com) to learn more.

INDICATION AND USAGE

VRAYLAR is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about any medical conditions and if you (continued):

- are pregnant or plan to become pregnant. VRAYLAR may harm your unborn baby. Taking VRAYLAR during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth. Talk to your healthcare provider about the risk to your unborn baby if you take VRAYLAR during pregnancy. If you become pregnant or think you are pregnant during treatment, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics at 1-866-961-2388 or <http://www.womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. It is not known if VRAYLAR passes into breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with VRAYLAR.

Tell your healthcare provider about all medicines that you take, including prescriptions, over-the-counter medicines, vitamins, and supplements. VRAYLAR may affect the way other medicines work, and other medicines may affect how VRAYLAR works. Do not start or stop any medicines while taking VRAYLAR without talking to your healthcare provider.

What are the most common side effects of VRAYLAR?

- The most common side effects include difficulty moving or slow movements, tremors, uncontrolled body movements, restlessness and feeling like you need to move around, sleepiness, nausea, vomiting, indigestion, constipation, feeling tired, trouble sleeping, increased appetite, and dizziness.

These are not all the possible side effects of VRAYLAR.

Please see Important Safety Information throughout and full [Prescribing Information](#), including [Boxed Warnings](#), and [Medication Guide](#).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit [AbbVie.com/myAbbVieAssist](https://www.abbvie.com/myAbbVieAssist) to learn more